

JOURNAL OF NEMATOLOGY

Official Journal of

THE SOCIETY OF NEMATOLOGISTS

VOLUME 15

Quarterly in

1983

Editor-in-Chief

JOSEPH A. VEECH

Editorial Board

William W. Carter
Joseph A. Veech
A. C. Triantaphyllou

A. W. Johnson
James G. Baldwin
Lorin Krusberg

Associate Editors

A. F. Bird
E. Cohn
B. R. Kerry
J. H. O'Bannon

D. P. Schmitt
R. A. Rohde
Stephen A. Lewis
Randy Gaugler

Technical Editor

Kathleen K. Leabo

CONTENTS

JOURNAL OF NEMATOLOGY

Vol. 15, No 1

January 1983

| | |
|---|-----|
| An Illustrated Key to the Cyst-Forming Genera and Species of Heteroderidae in the Western Hemisphere with Species Morphometrics and Distribution: R. H. MULVEY and A. MORGAN GOLDEN | 1 |
| Redescription of <i>Heterodea zeae</i> , the Corn Cyst Nematode, with SEM Observations: A. MORGAN GOLDEN and R. H. MULVEY | 60 |
| Description of <i>Distolabrellus veechi</i> n. gen., n. sp. (Nematoda: Rhabditidae): R. V. ANDERSON | 70 |
| Description of <i>Paktylenchus tuberosus</i> gen. n., sp. n. (Nematoda: Tylenchinae) from Pakistan: M. A. MAQBOOL | 76 |
| Relationship Between <i>Paratrichodorus</i> sp. Density, and Growth of Wheat in Pots: R. N. INSERRA, M. DiVITO, N. VOVLAS and J. W. SEINHORST | 79 |
| Reproduction of <i>Criconemoides simile</i> , <i>Helicotylenchus pseudorobustus</i> , and <i>Paratylenchus projectus</i> on Soybean: E. C. McGAWLEY and R. A. CHAPMAN | 87 |
| Efficacy of Selected Nonvolatile Nematicides on Control of <i>Ditylenchus destructor</i> in Iris: WILLIAM A. HAGLUND | 92 |
| Effect of Time of Application on the Action of Foliar Sprays of Oxamyl on <i>Meloidogyne hapla</i> in Tomato: Z. A. STEPHAN and D. L. TRUDGILL | 96 |
| Development of <i>Meloidogyne incognita</i> Inhibited by <i>Digitaria decumbens</i> cv. Pangola: SANAA HAROON and GROVER C. SMART, JR. | 102 |
| Population Development and Reproduction of Fungus- and Bacterium-Feeding Nematodes in the Presence of Insect Growth Regulators: J. L. TOWNSHEND, D. J. PREE, and A. B. BROADBENT | 105 |
| Effect of Soil Water Potential on Survival of <i>Meloidogyne javanica</i> in Fallow Soil: A. J. TOWSON and W. J. APT | 110 |
| Life Cycle of <i>Heterodera zeae</i> Koshy, Swarup, and Sethi on <i>Zea mays</i> L. Axenic Root Explants: J. A. LAURITIS, R. V. REBOIS, and L. S. GRANNEY .. | 115 |
| Response of Soybean Lines to Differentially Selected Cultures of Soybean Cyst Nematode <i>Heterodera glycines</i> Ichinohe: S. C. ANAND and G. S. BRAR | 120 |
| Effects of Temperature and Root Leachates on Embryogenic Development and Hatching of <i>Meloidogyne chitwoodi</i> and <i>M. hapla</i> : R. N. INSERRA, G. D. GRIFFIN, and D. V. SISSON | 123 |
| Ultrastructural Changes Caused by <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> in <i>Meloidogyne javanica</i> Induced Giant Cells in <i>Fusarium</i> Resistant and Susceptible Tomato Cultivars: F. FATTAH and J. M. WEBSTER | 128 |
| Host Suitability and Susceptibility of <i>Glycine max</i> cv. Bedford to Race 1 of <i>Heterodera glycines</i> : J. L. STARR, D. P. SCHMITT, and A. W. DUPREE, JR. .. | 136 |
| RESEARCH NOTES | |
| Comparative Response to Sugarbeet and Fodder Beet to <i>Heterodera schachtii</i> : G. D. GRIFFIN and J. C. THEURER | 140 |

| | |
|---|-----|
| An Improved Technique for Clearing and Staining Plant Tissues for Detection of Nematodes: D. W. BYRD, JR., T. KIRKPATRICK, and K. R. BARKER | 142 |
| Reaction of <i>Beta</i> spp. to Root-knot Nematodes: M. DI VIRO | 144 |
| Observations on the Inner Labial Sensilla of Second-stage Larvae of <i>Heterodera glycines</i> : GREGORY R. NOEL and BARBARA A. STANGER | 146 |
| Gall Formation on <i>Pistacia vera</i> by <i>Rotylenchulus macrodorum</i> : N. VOVLAS | 148 |

Vol. 15, No. 2

April 1983

| | |
|---|-----|
| Nematode Chemosensilla: Form and Function: K. A. Wright | 151 |
| Biochemical Studies of Olfaction: Role of Cilia in Odorant Recognition: L. D. Rhein | 159 |
| Chemotactic Behavior of Nematodes: David B. Dusenbery | 168 |
| Hypotheses and Possibilities of Intervention in Nematode Chemoresponses: Bert M. Zuckerman | 173 |
| <i>Cryphodera utahensis</i> n. sp., (Heteroderidae), A New Species from Wild Rose in Utah: J. G. Baldwin, M. Mundo-Ocampo, and A. A. Othman | 182 |
| Descriptions of <i>Chitwoodius brasiliensis</i> n.sp., <i>Chitwoodius rusticulus</i> n.sp., and <i>Vanderlindia venata</i> n.sp. (Dorylaimida: Tylencholaimidae: Vanderlindiinae): Mohammad Rafiq Siddiqi | 192 |
| Effect and Reproduction of <i>Rotylenchulus reniformis</i> on Sweet Potato Selections: C. A. Clark and V. L. Wright | 198 |
| Population Dynamics of <i>Meloidogyne incognita</i> and <i>Rotylenchulus reniformis</i> Alone and in Combination, and Their Effects on Sweet Potato: Ronald J. Thomas and Christopher A. Clark | 204 |
| Parasitism of <i>Rotylenchulus reniformis</i> on Soybean Root <i>Rhizobium</i> Nodules in Venezuela: J. A. Meredith, R. N. Inserra, D. Monzon de Fernandez | 211 |
| Effects of Concomitant Development on Reproduction of <i>Meloidogyne incognita</i> and <i>Rotylenchulus reniformis</i> on Sweet Potato: Ronald J. Thomas and Christopher A. Clark | 215 |
| Interactions Between Nematodes and Other Factors on Plants: H. R. Wallace | 221 |
| Validation of a Model for Prediction of Host Damage by Two Nematode Species: Larry W. Duncan and Howard Ferris | 227 |
| Host-Parasite Relationships of <i>Atalodera</i> spp. (Heteroderidae): M. Mundo-Ocampo and J. G. Baldwin | 234 |

| | |
|--|-----|
| Population Development of <i>Pratylenchus hexincisus</i> in Eight Corn Inbreds: Laura Georgi, J. M. Ferris, and V. R. Ferris | 243 |
| Influence of Planting Date on Damage to Soybean Caused by <i>Heterodera glycines</i> : R. S. Hussey and H. R. Boerma | 253 |
| Host Response to <i>Sarisodera hydrophila</i> Wouts and Sher, 1971: M. Mundo-Ocampo and J. G. Baldwin | 259 |
| Effect of <i>Sesbania rostrata</i> on <i>Hirschmanniella oryzae</i> in Flooded Rice: G. Germani, G. Reversat, and M. Luc | 269 |
| Development of <i>Heterodera glycines</i> Ichinohe on Soybean, <i>Glycine max</i> (L.) Merr., under Gnotobiotic Conditions: J. A. Lauritis, R. V. Rebois, and L. S. Graney | 272 |
| Susceptibility of plant selections to <i>Heterodera schachtii</i> and a race of <i>H. trifolii</i> parasitic on Sugarbeet in The Netherlands: A. E. Steele, H. Toxopeus, and W. Heijbroek | 281 |
| Development of the False Root-knot Nematode, <i>Nacobbus aberrans</i> , on Sugarbeet: R. N. Inserra, N. Vovlas, G. D. Griffin, and J. L. Anderson | 288 |
| Development of <i>Thelastoma bulhoesi</i> (Oxyurata: Thelastomatida) and the Effect of Thiabendazole on the Unembryonated Egg: Gary L. McCallister and Gerald D. Schmidt | 296 |
| Parasitic Variability of <i>Meloidogyne incognita</i> Populations on Susceptible and Resistant Cotton: T. L. Kirkpatrick and J. N. Sasser | 302 |
| Effect of Soil Texture on the Distribution and Infectivity of <i>Neoaplectana carpocapsae</i> (Nematoda: Steinernematidae): Ramon Georgis and George O. Poinar, Jr. | 308 |
| Nematode parasites and associates of <i>Dendroctonus</i> spp. and <i>Trypodendron lineatum</i> (Coleoptera: Scolytidae), with a description of <i>Bursaphelenchus varicauda</i> n.sp.: Cyril H. S. Thong and John M. Webster | 312 |
| Plant-Parasitic Nematodes of South Viet Nam: Gguyen B. Khuong | 319 |
| <i>Hypoaspis</i> nr. <i>aculeifer</i> : a Mite Predacious on Root-knot and Cyst Nematodes: R. N. Inserra and D. W. Davis | 324 |
| Wall Structure of the Mature Female and Cystoid Body of <i>Meloidoderita kirjanovae</i> (Nematoda: Criconematoidea): E. Cohn and M. Mordechai | 325 |

Vol. 15, No. 3

July 1983

| | |
|---|-----|
| Effect of Soil Texture on the Distribution and Infectivity of <i>Neoaplectana glaseri</i> (Nematoda:Steinernematidae): Ramon Georgis and George O. Poinar, Jr. | 329 |
| Studies on Anhydrobiosis of <i>Pratylenchus thornei</i> : I. Glazer and D. Orion | 333 |

| | |
|---|-----|
| Biochemical Identification of the Two Races of <i>Radopholus similis</i> by Starch Gel Electrophoresis: R. N. Huettel, D. W. Dickson, and D. T. Kaplan | 338 |
| Biochemical Identification of the Two Races of <i>Radopholus similis</i> by Polyacrylamide Gel Electrophoresis: R. N. Huettel, D. W. Dickson, and D. T. Kaplan | 345 |
| <i>Meloidogyne aquatilis</i> n. sp. (Nematoda:Meloidogynidae) from <i>Spartina pectinata</i> with a Key to the Canadian Species of <i>Meloidogyne</i> : B. A. Ebsary and E. S. Eveleigh | 349 |
| Description of <i>Pararotylenchus belli</i> n.sp. (Nematoda:Hoplolaimidae): R. T. Robbins | 353 |
| Description of <i>Thecavermiculatus andinus</i> n.sp. (Meloidoderidae), a Round Cystoid Nematode from the Andes Mountains of Peru: A. M. Golden, J. Franco, P. Jatala, and E. Astogaza | 357 |
| <i>Pungentus porosus</i> n.sp. (Nematoda:Dorylaimoidea) from Canada with a Key to the Species of <i>Pungentus</i> : B. A. Ebsary | 363 |
| Morphological Characteristics of <i>Hoplolaimus indicus</i> Sher, 1963 in Canada, a Parasite of Wild Rice: R. V. Anderson | 366 |
| Fine structure of body wall cuticle of females of <i>Meloidodera charis</i> , <i>Atalodera lonicerae</i> , and <i>Sarisodera hydrophila</i> (Heteroderidae): J. G. Baldwin | 370 |
| <i>Meloidogyne enterolobii</i> n. sp. (Meloidogynidae), a Root-knot Nematode Parasitizing Pacara Earpod Tree in China: Baojun Yang and J. D. Eisenback | 381 |
| Histopathology of Selected cultivars of Tobacco infected with <i>Meloidogyne</i> species: C. Sosa-Moss, K. R. Barker, and M. E. Daykin | 392 |
| Influence of Maize Rotations on the Yield of Soybean Grown in <i>Meloidogyne incognita</i> Infested Soil: Robert A. Kinloch | 398 |
| Resistance of Some <i>Vitis</i> Rootstocks to <i>Xiphinema index</i> : A. R. Harris | 405 |
| Interaction of Vesicular-arbuscular Mycorrhizal Fungi and Phosphorus with <i>Meloidogyne incognita</i> on Tomato: K. M. Thomson Cason, R. S. Hussey, and R. W. Roncadori | 410 |
| Dispersion and Distribution of <i>Pratylenchus scribneri</i> and <i>Hoplolaimus galeatus</i> in Soybean Fields: T. Alby, J. M. Ferris, and V. R. Ferris | 418 |
| The Interrelationship of <i>Heterodera schachtii</i> and <i>Ditylenchus dipsaci</i> on Sugarbeet: G. D. Griffin | 426 |
| Population Dynamics of <i>Heterodera glycines</i> and Soybean Response in Soils Treated with Selected Nematicides and Herbicides: D. P. Schmitt, F. T. Corbin, and L. A. Nelson | 432 |
| On the Methodology of Nematode Extraction from Field Samples: Baermann Funnel Modifications: D. R. Viglierchio and Richard V. Schmitt .. | 438 |
| On the Methodology of Nematode Extraction from Field Samples: Density Flotation Techniques: David R. Viglierchio and Tom T. Yamashita | 444 |

| | |
|--|-----|
| On the Methodology of Nematode Extraction from Field Samples: Comparison of Methods for Soil Extraction: David R. Viglierchio and Richard V. Schmitt | 450 |
| Influence of Metalaxyl on Three Nematodes of Citrus: David T. Kaplan | 454 |
| <i>Rhizonema sequoiae</i> n.gen. n.sp. from Coast Redwood <i>Sequoia sempervirens</i> (D.Don) Endl.: I. Cid Del Prado Vera, B. F. Lownsbery and A. R. Maggenti | 460 |
| Effects of Selected Nematicides on Hatching of <i>Heterodera schachtii</i> : Arnold E. Steele | 467 |

Vol. 15, No. 4

October 1983

| | |
|--|-----|
| Abstracts of Papers Presented at the Twenty-Second Annual Meeting of the Society of Nematologists, Ames, Iowa, 26-30 June 1983 | 475 |
| Control of <i>Globodera rostochiensis</i> in Relation to Method of Applying Nematodes: B. B. BRODIE | 491 |
| Effects of Dosage Sequence on the Efficacy of Nonfumigant Nematicides, Plantain Yields, and Nematode Seasonal Fluctuations as influenced by Rainfall: T. BADRA AND F. E. CAVENESS | 496 |
| Use of Avermectins for the Control of <i>Meloidogyne incognita</i> on tomatoes: SAHAG GARABEDIAN AND S. D. VAN GUNDY | 503 |
| Effect of Aldicarb, Ethoprop, and Carbofuran on Control of Coffee Root-knot Nematode, <i>Meloidogyne exigua</i> : S. P. HUANG, I. C. RESENDE, D. E. DE SOUZA, AND V. P. CAMPOS | 510 |
| Parasitic Habits of <i>Trophotylenchulus floridensis</i> (Tylenchulidae) and its Taxonomic Relationship to <i>Tylenchulus semipenetrans</i> and Allied Species: E. COHN AND D. T. KAPLAN | 514 |
| Studies on <i>Lasioseius scapulatus</i> , a Mesostigmatid mite predaceous on nematodes: I. IMBRIANI AND R. MANKAU | 523 |
| Electron Microscope Characterization of Carbohydrate Residues on the Body Wall of <i>Xiphinema index</i> : Y. SPIEGAL, W. M. ROBERTSON, S. HIMMELHOCH AND B. M. ZUCKERMAN | 528 |
| <i>Caenorhabditis elegans</i> : Stage Specific Differences in Cuticle Surface Carbohydrates: BERT M. ZUCKERMAN AND ITZHAK KAHANE | 535 |
| Interrelationships between Ethylene Production, Gall Formation, and Root-knot Nematode Development in Tomato Plants Infected with <i>Meloidogyne javanica</i> : I. GLAZER, D. ORION, AND A. APELBAUM | 539 |
| Host Response to <i>Meloidodera</i> spp. (Heteroderidae): M. MUNDO-OCAMPO AND J. G. BALDWIN | 544 |
| Effect of Plant Age and Transplanting Damage on Sugar Beets infected by <i>Heterodera schachtii</i> : TH. H. A. OLTJOF | 555 |
| Influence of Six Vegetable Cultivars on Reproduction of <i>Meloidogyne javanica</i> : N. D. BAFOKUZARA | 559 |

| | |
|---|-----|
| Influence of <i>Meloidogyne hapla</i> Chitwood, 1949 on Development and Establishment of <i>Heterodera schachtii</i> Schmidt, 1871 on <i>Beta vulgaris</i> L: PARVIZ JATALA AND HAROLD J. JENSEN | 564 |
| Effect of an Extract from Saprozoic Nematode-Infested Compost on the Mycelial Growth of <i>Agaricus brunnescens</i> : T. D. KAUFMAN, J. R. BLOOM, AND F. L. LUKEZIC | 567 |
| Biology of <i>Heterodera mediterranea</i> : N. VOVLAS AND R. N. INSERRA | 571 |
| Reproductive Development of <i>Verutus volvingentis</i> (Tylenchida: Heteroderidae): R. P. ESSER | 576 |
| Taxonomic Limits of the Genus <i>Nagelus</i> (Thorne and Malek, 1968) Siddiqi, 1979 with a Description of <i>Nagelus borealis</i> n.sp. from Alaska: THOMAS O. POWERS, JAMES G. BALDWIN, AND A. H. BELL | 582 |
| Effect of <i>Aedes aegypti</i> (Diptera: Culicidae) Age on Sex Ratios in <i>Romanomermis culicivora</i> (Nematoda: Mermithidae): NAYNA VYAS-PATEL | 594 |
| Susceptibility of the Colorado Potato Beetle and the Sugarbeet Wireworm to <i>Steinernema feltiae</i> and <i>S. glaseri</i> : H. HAROLD TOBA, JAMES E. LINDEGREN, JOHN E. TURNER, AND PATRICK V. VAIL | 597 |
| Characteristics of Herbicide Distribution as Patterns for Nematicide Behavior: D. E. BAYER | 602 |
| Progress and Prospects in the Chemotherapy of Nematode Infections of Man and Other Animals: W. C. CAMPBELL | 608 |
| Insecticides and Their Design: HANS E. HUMMEL | 615 |
| RESEARCH NOTES | |
| Reproduction of Two Races of <i>Meloidogyne incognita</i> in Tomato Plants Grown at High Temperature: M. T. ARAUJO, D. W. DICKSON, J. J. AUGUSTINE, AND M. J. BASSET | 640 |
| <i>Monacrosporium lysipagum</i> Infecting Egg Masses of <i>Meloidogyne acrita</i> : R. P. ESSER | 642 |
| Response of Two Alfalfa Cultivars to <i>Meloidogyne hapla</i> : R. N. INSERRA, J. H. O'BANNON, M. DI VITO, AND H. FERRIS | 644 |
| Root Extracts of Pangola Digitgrass Affect Egg Hatch and Larval Survival of <i>Meloidogyne incognita</i> : SANAA HAROON AND GROVER C. SMART, JR. ... | 646 |
| Effects of Pangola Digitgrass on <i>Meloidogyne arenaria</i> , <i>M. javanica</i> , and <i>M. hapla</i> : SANNA HAROON AND GROVER C. SMART, JR. | 649 |
| Isolation of an Unusual Entaphelenchid (Tylenchida: Entaphelenchidae) from the Hemocoel of <i>Platysoma punctigerum</i> LeConte Coleoptera: Histeridae): ROBIN M. GIBLIN AND HARRY K. KAYA | 650 |
| Vertical Migration of <i>Heterorhabditis bacteriophora</i> and <i>H. heliothidis</i> (Nematoda: Heterorhabditidae) in Sandy Loam Soil: RAMON GEORGIS AND GEORGE O. POINAR, JR. | 652 |

Information for Contributors

Original papers on basic, applied, descriptive, or experimental nematology are considered for publication. Other categories include research notes (not longer than four pages of copy, including tables and figures); reviews developing new concepts, hypotheses, or theories; and abstracts of papers presented at the annual meetings. At least one author must be a member of the Society of Nematologists, except that nonmember staff of subscribing institutions outside the United States or Canada may submit one paper per volume.

Manuscripts: All manuscripts will be in English, although abstracts may be in a "congress" language other than English. All copy, including tables, figure legends, and literature citations, should be typed with double spacing on 8 1/2 × 11-inch manuscript paper with wide margins all around; words should not be broken (hyphenated) at the end of lines. Authors should have the manuscript critically reviewed by two colleagues whose names should be included in the letter of submission. Three copies (the original and two Xerox or similar quality copies) should be sent to the editor-in-chief with the understanding that publication is to be exclusively in the JOURNAL OF NEMATOLOGY. Papers will be reviewed by specialists in the field and edited by selected editorial staff members.

Address: Dr. J. A. Veech, Editor-in-Chief, JOURNAL OF NEMATOLOGY, National Cotton Pathology Research Laboratory, P. O. Box JF, College Station, TX 77840.

Style: The CBE Style Manual, Fourth Edition, and Webster's Third New International Dictionary will be guides for form, style, and spelling. Use SI (Système International) units of measurement; if British and Fahrenheit are also desired, include them in parentheses. Author(s) name(s) and page number should appear in the upper right corner of each page. The title page should be numbered 1, the abstract page 2, the first page of the text 3, etc.

Title page: Should contain the title, name(s) of author(s), a hairline over the notation "Received for publication _____"; footnote(s) giving institutional affiliation(s), senior author's address (may be the same), and acknowledgments (if any) and, finally, a condensed running head of no more than 60 spaces (including author's last name). The title should utilize strong key words and be brief, preferably not more than 100 spaces. Scientific names of plants and animals should be given. Trade names should not be used in the title.

Abstract page: Should contain an abstract—no more than 1,300 spaces—and key words *not* appearing in the title. (See Journal for style.) The abstract serves as the summary.

Body: The body of the manuscript should be organized after the introduction (do not use the word "Introduction") according to "Materials and Methods," "Results," "Conclusions," "Discussion," "Literature Cited," "Tables," "Legends for Figures," and "Figures."

Literature citations: Cite by number in parentheses in the text. Under "Literature Cited," list (double spaced, no underlining) on a separate page alphabetically by senior author and numbered consecutively. Each citation should be checked against the original publication to verify author's name,

year, correct title, journal name (or publisher name and location), volume number, and inclusive paging. Journal names will be abbreviated as in the "List of Serials" in *Biological Abstracts*.

Tables: Number (Arabic) each table and place on a separate page with descriptive title at the top. Data will appear in Arabic numerals with minimum footnotes identified by symbols as indicated in the CBE Style Manual. Vertical lines must be avoided. Table dimensions should consider printed page size limitations.

Figures: Number (Arabic) all "Legends for Figures" and type consecutively on a single page. In planning large illustrations, remember that the printed page is 140 mm wide × 213 mm deep. Make allowances for the legend. Combine and mount several illustrations into composite plates when possible. Trim waste from figures and leave a space of 1–2 mm between photographs in a composite. Do not send original illustrations of taxonomic papers until requested, but provide a complete set of figures (photocopies okay) with each copy of each manuscript. Lettering on illustrations should be of high contrast, neat, and at least 1 mm high after final reduction. Identify each figure on the back lightly with a soft pencil—with "J. Nematol.," manuscript number, author's name, figure number, and width desired. Color illustrations and folded inserts can be printed only if the extra costs are paid for by the author.

Research notes: Same as for Manuscripts except there will be no abstract page. Key words should be included, however.

Biological nomenclature: Authors should follow the "International Rules of Nomenclature" in naming organisms; each should be identified by genus, species, authority (if applicable), varietal names (in single quotes), breeds, races, and accession number. Complete authorities and dates are required in taxonomic papers. Binomials should be underlined in typewritten copy (but not in "Lit. Cited") list and fully spelled out the first time they are used either in text, figure captions, or table titles. When new names of nematodes are introduced, the recommendations of the "International Code of Zoological Nomenclature," 1961, should be strictly applied, especially regarding type designation and statements indicating where such type material is deposited.

Chemical nomenclature: Use approved common (generic) names where available; otherwise use capitalized trade names followed by their chemical names. State the concentration of active ingredient. Chemical doses should be reported as amount of active or technical material per unit area, volume, or weight (e.g., 10 ml/1,000 m², 10 µg/ml, or 1 mg/kg). The chemical formulation should be given and method of application clearly stated.

Editorial page charge: In order to partially cover the cost of publishing manuscripts in this journal the Society suggests a contribution of \$35 per printed page, as it is impossible to meet the demand for journal space without some financial assistance. Payment is not a condition of publication, however, since articles are accepted on the basis of merit.

Separates: No free copies. Quotation and order blanks for reprints will be sent with proofs. Plate of figures are available upon request.